

CLINICAL HANDBOOKS BY DON MEICHENBAUM

TREATMENT OF INDIVIDUALS WITH ANGER-CONTROL PROBLEMS AND AGGRESSIVE BEHAVIOR

PRICE \$55 US Funds

TABLE OF CONTENTS (446 Pages – Softcover)

BY WAY OF INTRODUCTION

SECTION I: THE NATURE OF ANGER

SECTION II: A DEVELOPMENTAL LIFE-SPAN PERSPECTIVE OF AGGRESSIVE BEHAVIOR

SECTION III: ASSESSMENT

SECTION IV: INTRODUCTION TO TREATMENT

SECTION V: INITIAL PHASE OF COGNITIVE BEHAVIORAL TREATMENT

SECTION VI: SKILLS TRAINING PHASE

SECTION VII: GENERALIZATION GUIDELINES AND APPLICATION TRAINING

SECTION VIII: REFERENCES AND APPENDICES

ASSESSING AND TREATING ADULTS WITH PTSD

PRICE \$65 US Funds

600 Pages Softcover

For orders:

Make check payable to Don Meichenbaum

(No credit cards accepted) Price includes all costs (shipping and handling).

**Please mail check and orders to: Dr. Donald Meichenbaum
1560 Gulf Blvd.
Apt. 1002
Clearwater, FL 33767**

Email: dmeich@watarts.uwaterloo.ca

**SPECIAL OFFER—BOTH HANDBOOKS FOR
\$100---A SAVINGS OF \$20**

TREATMENT OF INDIVIDUALS WITH ANGER-CONTROL PROBLEMS AND AGGRESSIVE BEHAVIORS: A CLINICAL HANDBOOK

HOW TO USE THIS HANDBOOK	13
BY WAY OF INTRODUCTION	14
SECTION I: THE NATURE OF ANGER	18
SECTION II: A DEVELOPMENTAL LIFE-SPAN PERSPECTIVE OF AGGRESSIVE BEHAVIOR	48
SECTION III: ASSESSMENT	132
SECTION IV: INTRODUCTION TO TREATMENT	198
SECTION V: INITIAL PHASE TO COGNITIVE BEHAVIORAL TREATMENT	217
SECTION VI: SKILLS TRAINING PHASE	271
SECTION VII: GENERALIZATION GUIDELINES AND APPLICATION TRAINING	333
SECTION VIII: REFERENCES, APPENDICES AND WEBSITES	365

HOW TO USE THE ANGER-CONTROL HANDBOOK (User-friendly Guide)

I WANT TO LEARN MORE ABOUT OR HOW TO:

Conduct APPLICATION TRAINING (p. 341)
 Conduct ASSESSMENT (p. 132)
 Conduct ATTRIBUTION TRAINING (p. 347)
 Formulate CASE CONCEPTUALIZATION MODEL.
 (p. 82)
 Examples: Intimate Partner Violence (p. 108)
 Juvenile Offenders (p. 83)
 "DEFUSE" ANGRY INDIVIDUALS (p. 400)
 Consider EPIDEMIOLOGY (p. 26)
 Conduct COGNITIVE RESTRUCTURING
 ("RETHINKING" PROCEDURES) (p. 299)
 EDUCATE PATIENTS (p. 250)
 Identify CORE TASKS OF THERAPY (p. 204)
 Train FRONTLINE STAFF (p. 192)
 Understand GENDER DIFFERENCES (p. 71)
 Foster GENERALIZATION (p. 334)
 Employ GESTALT-BASED ASSESSMENT (p. 178)
 Conduct GOAL ATTAINMENT SCALING (p. 238)
 Conduct GOAL SETTING (p. 227)
 Conduct INTERVIEWS (p. 147)
 Examples: Clinical Interviews (p. 150)
 Life Space Crisis Interviews (p. 166)
 Use IMAGERY-BASED ASSESSMENT (p. 167)
 Employ IMAGERY-BASED REHEARSAL (p. 341)
 Employ LETTER WRITING/JOURNALING (p. 309)
 How to MAKE A BATTERER (p. 118)
 JUVENILE OFFENDER (p. 54)
 TERRORIST (p. 130)

Examine META-ANALYSES
 Child (p. 201)
 Adult (p. 199)
 Examine PATIENT HANDOUTS (p. 12)
 Consider PREVENTATIVE PROGRAMS (p. 122)
 Code PROVOCATIONS (p. 264)
 Assess PATIENT SATISFACTION (p. 354)
 Conduct PROBLEM-SOLVING TRAINING (p. 311)
 Conduct RELAPSE PREVENTION (p. 355)
 Conduct RELAXATION TRAINING (p. 278)
 Identify RESILIENCE FACTORS (p. 77)
 Conduct ROLE PLAYING (p. 329)
 Engage patients in SELF-MONITORING (p. 174)
 Examples: Anger Logs (p. 175)
 Conduct SKILLS TRAINING
 Examples: Communication Skills (p. 317)
 Interpersonal Skills (p. 324)
 Self-management Skills (p. 324)
 Employ SOCIAL INFORMATION PROCESSING
 MODEL (p. 35)
 Conduct SELF-INSTRUCTIONAL TRAINING (p. 284)
 Enhance THERAPEUTIC ALLIANCE (p. 218)
 Teach TIME-OUT PROCEDURES (p. 275)
 Address TREATMENT NONADHERENCE (p. 203)
 Predict VIOLENCE (DANGEROUSNESS) (p. 191)
 Examples: Intimate Partner Violence (p. 118)
 Psychiatric Patients (p. 179)
 Juvenile Offenders (p. 54)
 School Shooters (p. 15)
 Lookup WEBSITES (p. 435)

ASSESSING AND TREATING ADULTS WITH PTSD: A CLINICAL HANDBOOK

Donald Meichenbaum, Ph. D.

600 Pages – Softcover

-
- | | |
|-------------|--|
| Section I | <ul style="list-style-type: none">-- Epidemiological And Diagnostic Information-- Consider the nature and impact of natural, technological, and human-made disasters as evident in specific "victim populations"-- Critique diagnostic alternatives and "stage" theories |
| Section II | <ul style="list-style-type: none">-- Conceptualization of PTSD-- Reviews alternative conceptualizations and offers a "constructive narrative perspective" |
| Section III | <ul style="list-style-type: none">-- Assessment of PTSD-- Comprehensive enumeration of PTSD and related measures of comorbidity-- Describes a sequential gating assessment strategy-- Considers potential "positive" effects-- Includes the "best" clinical questions you can ask |
| Section IV | <ul style="list-style-type: none">-- Cautions About Assessment-- Consider the controversy over so-called "false memories"-- How to help the helpers |
| Section V | <ul style="list-style-type: none">-- Treatment Alternatives: A Critical Analysis-- Critically evaluates pharmacological, exposure, eye-movement desensitization, group interventions and other procedures-- Provides treatment guidelines and considers factors that influence the length of treatment |
| Section VI | <ul style="list-style-type: none">-- Specific Treatment Procedures: Practical "How to" Guidelines-- "How to": Educate clients about PTSD; deal with flashbacks; intrusive ideation; guilt; anger; addictive behaviors; depression; anxiety; conduct "memory work"; and address issues of multiple and borderline personality disorders-- Techniques include Stress inoculation training, cognitive restructuring, problem-solving, relapse prevention and family-based interventions |
| Section VII | <ul style="list-style-type: none">-- Post Disaster Interventions-- Consider who is most at "high risk"-- Describes and critiques Critical Incident Stress Debriefing; When can CISD make individuals "worse"?-- Workplace, accident, community interventions-- Consider the role of religion and rituals-- Over 1500 references |
-

For further information please call:
(519) 885-1211, extension 2551
or email: dmeich@watarts.uwaterloo.ca

TREATMENT CHALLENGES: FACT SHEET

(American Psychiatric Association, 1995; Institute of Medicine, 1990; McCrady & Ziedonis, 2001; Miller et al., 1995; Morgenstern et al., 2001; Najavits, & Weiss, 1994; Vaillant, 1983; Wampold et al., 1997; Zucker, 1994)

- The lifetime prevalence of drug dependence is 9.2% males and 5.9% females in the U.S., as reported in the National Comorbidity Study (Warner et al., 1995).
- Estimates of the number of children in the U.S. population having at least one parent who abuses alcohol or drugs range from 7-10 million. SUDS parents are 3 times more likely to neglect their children than control parents. Among children reported to Child Protection Services for various forms of abuse, 80% have parents with SUDS (Dunn et al., 2002).
- Individuals with addictive disorders represent an **heterogeneous population** with different etiologies and diverse courses. 80% of alcoholics never seek treatment with self-help or with treatment programs (Approximately, 10 million individuals with SUDS in U.S. do not seek treatment)
- Lifetime prevalence for alcohol abuse, alcohol dependence, or both is approximately 24% for males and 5% for females. Women comprise about 1/3 of those with substance abuse disorders.
- **Relapse rates** across chemical addictions (heroin, cocaine, nicotine, alcohol) and across various treatment modalities are **fairly uniform** and **discouraging** – around 75%. The likelihood of life-long abstinence is low.
- Among alcoholics who have been treated
 - ❖ 1/2 will be abstinent at 3 months
 - ❖ 1/3 will be abstinent at 6 months
 - ❖ 1/8 will be abstinent at 12 months
 - ❖ 1/10 will be abstinent at 18 months
- Approximately 90% of treated alcoholics have had at least one drink within 3 months of abstinence treatment. 45% - 50% will return to pretreatment drinking levels within a year.
- Overall, about 20% to 30% of alcoholic patients evidence long-term success with traditional treatments.

Meichenbaum

- 70% of those who **relapse**, will do so during the **first 3 months** after discharge. Nearly all those who relapse do so before 6 months expire. The first 90 days is the most vulnerable period for relapse across various substances of abuse (heroin, smoking, alcohol).
- Over 50% of those who enter treatment will **drop out within the first month**. Those who drop out of treatment have worse outcomes. 70% of those who complete aftercare vs. 23% of those who dropout are abstinent after 9 months. For example, only 54% of subjects completed treatment in PROJECT MATCH and only 27% completed treatment in Morgenstern et al. (2001) major treatment community study.
- Among those seeking consultation about 20% **will abstain without formal treatment** and an additional 21% **will moderate their drinking**.
- An **intensive inpatient treatment** program is **no more effective** than less intense treatment in outpatient settings.
- Studies that have compared differing lengths of treatment for alcohol use have not found differential positive effects for longer lengths of treatment. Increasing the length and intensity of treatment may be more important in treating patients with more severe dependence and co-occurring psychiatric problems.
- Low intensity interventions that focus on assessment, feedback and recommendations to reduce heavy drinking can be effective in reducing high-risk drinking.
- There is a strong association between positive treatment outcome and aftercare attendance and treatment compliance.
- **Treatment attendance** and the **degree of therapeutic alliance** account of the largest effect and explain more variance than all other predictors combined. **There is a stronger relationship between nonspecific aspects of treatment and outcome than between active ingredients and outcome.**
- Those individuals with addictive behaviors who have **social supports for abstinence** and who are a part of social network who are involved in treatment have more favorable prognoses.
- Ample evidence demonstrates that the **treatment of additional presenting problems** leads to more positive treatment outcomes than attention to the substance use disorder alone.
- Cognitive behavioral treatment (CBT) has been found to be more effective as one component of intensive treatment programs than as a stand alone intervention. CBT places primary focus not on alcohol consumption per se, but on life areas related functionally to drinking and relapse. Research has not yet established why CBT is an effective treatment for alcohol dependence (Morgenstern & Longabaugh, 2000).

FACILITATING TREATMENT ADHERENCE

**Donald Meichenbaum, Ph.D.
Distinguished Professor Emeritus,
University of Waterloo
and
Research Director of
The Melissa Institute for Violence Prevention and
Treatment of Victims of Violence
Miami, Florida
(www.melissainstitute.org)**

Address: University of Waterloo
Department of Psychology
Waterloo, Ontario
N2L 3G1
Phone: (519) 885-1211 ext. 2551
Fax: (519) 746-8631
Email: dmeich@watarts.uwaterloo.ca

Florida Address
(Nov. – May)
1560 Gulf Blvd.
Apt. 1002
Clearwater, FL 33767

Meichenbaum

Predicting Outcome

Brief therapies have been found to be as effective as long-term therapy for a number of psychiatric disorders (but not BPD).

The pattern of patient-therapist relationship established within the first three sessions is predictive of treatment outcome^{6,7}.

60% to 80% of symptom reduction in cognitive-behavior therapy of depression occurs within the first four sessions⁸

There now exist a list of empirically validated therapies (EVT)⁹ and a list EVT treatment manuals^{10,11}.

Meichenbaum

ADDRESSING ISSUES OF TREATMENT NONADHERENCE

(See Meichenbaum & Turk, 1983; Meichenbaum & Fong, 1993)

Factors Related to Patient Noncompliance: Challenge

Understanding the Reasons for Treatment Nonadherence

Intervention Strategies

Develop a Therapeutic Alliance

Collaborative Goal-Setting

Use Motivational Interviewing

Employ Short-term Interventions